

Impacts of Habitat Restoration and the Status of Avian Communities in Seattle City Parks





© DOUG SCHURMAN

Heermann's Gull



© TOM SANDERS

Anna's Hummingbird



© DOUG SCHURMAN

Yellow Warbler



© DOUG SCHURMAN

Lesser Scaup



© TIM BOYER

Bottom left: A volunteer engaged in the Neighborhood Bird Project at Magnuson Park counts gulls and waterfowl on a platform in Lake Washington.

BACKGROUND

THE NEIGHBORHOOD BIRD PROJECT (NBP) is a citizen science initiative that began in 1994, conceived, developed and managed by the Seattle Audubon Society. The NBP has two main goals; the first to monitor trends in avian abundance in Seattle City parks and green spaces. The second aims to empower citizens in becoming advocates for birds and wildlife habitat in their neighborhoods and communities. Monthly surveys are conducted by teams of volunteer bird watchers who conduct surveys at eight King County parks and green spaces – Carkeek Park, Golden Gardens Park, Discovery Park, Seward Park, Genesee Park, Washington Park Arboretum, Magnuson Park and Lake Forest Park. The data from these surveys provide an insight into the avian diversity and abundance in urban areas and affords an appreciation of the diversity that can be found in cities given appropriate quality habitat.

Introduction

CITY PARKS PRESENT A UNIQUE OPPORTUNITY for public engagement with nature, as well as providing habitat for wildlife and ecosystem services for millions of urban residents. Within the urban core of Seattle, city parks have been the focus of many habitat restoration projects to improve habitat quality and restore degraded lands – efforts supported by community members, local government, and nonprofit groups with a shared interest in maintaining biodiversity and native habitats easily accessible to the public. Because of this widespread public interest and ease of access, city parks are excellent targets for involving members of the public in long-term biological monitoring efforts at a greater frequency or scale than is typically possible for sites in remote areas or for surveys conducted by professional scientists.

In order to monitor trends in avian diversity and abundance over time, and to take advantage of the expertise and enthusiasm of volunteers from the surrounding

communities, the Seattle Audubon Society started the Neighborhood Bird Project (NBP) in 1994, with a series of volunteer-led surveys in Carkeek Park. Surveys have since expanded to seven other sites, and today are conducted once a month, year-round, at each of over 200 survey points distributed in natural or restored habitats in the Seattle area. Here we present a summary of findings from the first 17 years of NBP surveys in four Seattle City Parks: Discovery Park, Golden Gardens, Carkeek Park, and Magnuson Park (see Appendix for maps). The primary goals of this analysis are (1) to summarize general trends in avian diversity and abundance over time in the study areas, and (2) to assess the impact of habitat restoration activities conducted in the vicinity of survey points on bird communities. We examined overall trends across all species and parks and a detailed assessment of observed differences between restored and non-restored sites.

Survey Methods and Focal Species/Groups

NBP point counts are conducted by teams of volunteers at eight city parks once each month. Point count stations are located at least 200m apart at pre-determined locations on walking loops, with each loop including 5-9 stations. Following an initial one-minute rest period after arriving at a point count station, surveyors record the species, number, and mode of detection (seen/heard/flyover) of any birds observed within 50m of the survey point in a 5-minute period. Surveyors also record a brief description of the weather and wind conditions at the time of the survey. Surveys were not conducted during very poor weather (snow, heavy rain or wind). Data collection began in Carkeek Park in 1997 and expanded to other sites through 2003 with the addition of Discovery Park.

Five groups of species were selected for focused analysis in this report in order to represent communities of particular

interest to biologists and land managers and to allow us to draw some conclusion about varying trends across taxa favoring different habitat types. We also selected six focal species for extra analysis, again to allow inference of trends in species of particular interest and to draw out patterns that are not apparent in a generalized analysis of diversity and abundance. Focal species were also selected to reflect divergent habitat preferences, in order to provide some assessment of the affect of restoration in different areas on different segments of the local bird community. Species groupings and focal species are summarized in Table 1.

Statistical analyses presented in this report were conducted in R (version 3.0.2) and visualized with the ggplot2 package. Geographic analyses and maps were prepared with ArcGIS (version 10.1, ESRI 2012).



© DOUG SCHUMMAN



© TOM SANDERS

Spotted Towhee

Downy Woodpecker

TABLE 1

Species groups and focal species.

GROUP	SPECIES	DESCRIPTION
Invasive Species	European Starling, Eurasian Collared-dove, House Sparrow	Non-native species known to displace natives, typically targeted for population reduction in restoration projects.
Human-associated Species	American Crow, Rock Pigeon, European Starling, House Sparrow	Common urban birds with high populations in disturbed areas.
Riparian Species	Orange-crowned Warbler, Wilson’s Warbler, Song Sparrow, Belted Kingfisher, Yellow Warbler, Common Yellowthroat	Species that nest or primarily inhabit brushy habitat adjacent to waterways. Typically targeted for population increase in restoration projects.
Warblers	Orange-crowned Warbler, Wilson’s Warbler, Black-throated Gray Warbler, Common Yellowthroat, Macgillivray’s Warbler, Hermit Warbler, Townsend’s Warbler, Yellow Warbler, Yellow-rumped Warbler	Colorful, vocal, long-distance migrants; including many of our most charismatic breeding-season taxa.
Woodpeckers	Hairy Woodpecker, Downy Woodpecker, Pileated Woodpecker, Northern Flicker	A diverse family with large populations in many parks – sensitive to a variety of habitat changes.
FOCAL SPECIES	PREFERRED HABITAT	NOTES
Anna’s Hummingbird, <i>Calypte anna</i>	Generalist	Documented local population increases suggest increasing availability of food and habitat, especially in winter.
Savannah Sparrow, <i>Passerculus sandwichensis</i>	Meadows, grasslands, and some shrub-steppe habitats in suitable areas.	A common breeder in open areas of Discovery and Magnuson Parks.
White-crowned Sparrow, <i>Zonotrichia leucophrys</i>	Meadows or grasslands with scattered shrubs, shrub-steppe.	Both wintering and resident populations present throughout the year.
Brown Creeper, <i>Certhia americana</i>	Mature coniferous forest.	A little-seen resident species that forages on the trunks of large conifers.
American Crow, <i>Corvus brachyrhynchos</i>	Generalist	Perhaps Seattle’s most successful species; abundant in many disturbed habitats.
Wilson’s Warbler, <i>Wilsonia pusilla</i>	Thick mid-succession understory growth or riparian thickets.	A common neotropical migrant and summer resident in suitable habitat.

Status and Trends in Avian Diversity in Seattle City Parks

NBP surveys recorded 232 species in Seattle City Parks over a 17-year timespan. Total species diversity (the number of species reported in a given park over the entire study period) is highest in Discovery Park with 207 species, followed by Magnuson Park (169), Golden Gardens (123), and Carkeek Park (114, Figure 1). Mean annual species diversity (the average number of species reported per year) is 130 species across all parks from 2003 to 2013, with individual parks ranked from Magnuson (87 species per year) to Golden Gardens (48 species per year).

Citizen science volunteers taking part in the Neighborhood Bird Project at Golden Gardens Park peer into the tree canopy to count birds during a survey.



To assess general trends in avian diversity and abundance across time, and to establish a baseline for continued long-term study of the impact of restoration activities, we focused on two measures: annual species diversity (the number of species recorded in a park in a given year) and mean abundance (the average number of birds recorded per station per survey; this can also be thought of as the frequency of occurrence). As a simple test of change over time, both measures were plotted by year and fit with a linear model. This allows us to infer relative rates of change of the populations of different species within the parks (see Figure 2). Separate models were applied before and after 2003 for focal species and species groups analyses, as the addition of the large number of survey points in Discovery Park that year introduced new habitat diversity into the dataset and makes direct comparisons with the abundance and diversity prior to 2003 unreliable for most species.

Our analyses of the NBP data found suggestive trends of decreasing species diversity through time in most locations surveyed – on the order of 1 fewer species per year across all parks – but this pattern was not strongly supported by linear models (Table 2, Figure 3). Of the four parks assessed, all but Discovery Park showed slightly decreasing species abundance, but none of the models explained more than 20% of the variation in the data, suggesting that documented trends in species diversity reflect random variation (or at least nonlinear change over time) rather than any consistent single pattern of change over the study period. Furthermore, the trend towards decreasing diversity over time disappears when the unusually low diversity numbers for 2012 – when fewer surveys were completed – are removed, suggesting that variation in survey effort is responsible for much of the

© TIM BOYER



© DOUG SCHUMMAN

Brown Creeper

SPECIES RICHNESS BY PARK




FIGURE 1

Number of species recorded in each park

AVERAGE RATE OF CHANGE

in Mean Per-survey Abundance (2003-2014)




FIGURE 2

Ranked slopes of linear regression lines for mean number of individuals detected per-survey, per-station. Limited to species with mean annual detections greater than 10 and mean rates greater than 0.0075 detections/station/survey/year.

TABLE 2

Trends in annual species diversity over time.


LOCATION	MEAN ANNUAL SPECIES DIVERSITY	TREND	SLOPE (SPECIES PER YEAR)	R ²
All Parks	130.000	Decreasing	-1.2	0.019
Discovery Park	78.889	Increasing	1.1	0.058
Magnuson Park	86.667	Decreasing	-0.65	0.19
Carkeek Park	54.389	Decreasing	-0.3	0.055
Golden Gardens Park	48.222	Decreasing	-0.32	0.009

SPECIES REPORTED

All Parks

FIGURE 3

Annual species diversity over time.





Savannah Sparrow

decline. Although any decline in species diversity is a cause for concern, discerning long-term decline from random (or at least unmeasured) variation is a difficult task even for the best-designed surveys, and to date the NBP data shows no strongly supported pattern of change. However, continued data collection under a comparable protocol will maximize the value of data already collected and may serve to point out important trends in species diversity in the future.

Focusing on species groups, the data suggest that invasive and human-associated species have decreased in relative abundance while woodpeckers and warblers have increased. Across all parks, the frequency of invasive birds has declined consistently since the start of surveys, decreasing from over 3 per survey per station in 1997 to fewer than 1 per survey per station across all sites surveyed to date in 2014. The frequency of occurrence of human-associated species has also declined, while frequency of riparian birds increased steadily prior to the inclusion of Discovery Park sites in 2003 and has since held steady. Both warblers and woodpeckers showed slight, steady increases in frequency of occurrence across all parks (Figure 4).

Among the focal species studied, Anna's Hummingbird, *Calypte anna*, and Brown Creeper, *Certhia americana*, showed the most consistent trends across all parks. Anna's Hummingbird increased in frequency roughly 50% from 2004 to 2013 (Figure 5) – an impressive rate of increase roughly in line with other observed increases in populations of this species throughout the northwest, likely driven by an increase in winter food availability from decorative plantings and hummingbird feeders. Brown Creeper also increased significantly throughout the study period, though higher

variation in counts of this species meant that the linear trend explained less of the total variation than was the case with the Anna's Hummingbird.

Although the two focal species selected to represent birds preferring open meadow habitats – White-crowned Sparrow, *Zonotrichia leucophrys* and Savannah Sparrow, *Passerculus sandwichensis*, – did not show any well-supported linear trends across all parks, local patterns of abundance were variable and deserve careful observation as restoration and maintenance work is ongoing. Savannah Sparrow in particular is rarely observed in either Golden Gardens or Carkeek parks, but breeds abundantly in both Magnuson and Discovery Parks. Since the beginning of data collection Savannah Sparrow frequency has nearly doubled in Magnuson Park, but has declined by roughly half in Discovery Park (Figure 6).

The cause of these local changes in Savannah Sparrow abundance is not directly addressed by these data, but differences in the timing and amount of restoration and land management activities between sites likely plays a role. As the timing of mowing in grassland habitats in Discovery Park and its impact on grassland-nesting birds has long been a point of contention between birders and land managers at the site, this study's observation of long-term decline in abundance should serve as a useful data point in calibrating further management actions in the area.

FIGURE 4 Species group trends in abundance.







FIGURE 5 Anna's Hummingbird and Brown Creeper change in detection frequency over the survey period.





FIGURE 6 Trends in Savannah Sparrow abundance in Discovery and Magnuson Parks.



Impacts of Habitat Restoration


Green Seattle Partnership (GSP) has been conducting both professional and volunteer-driven habitat restoration projects throughout the Seattle area since 2004 and currently organizes ongoing restoration and monitoring projects within 50m of the majority of survey points incorporated in the NBP dataset (see Appendix). GSP restoration projects typically proceed through four phases: 1 – invasive removal, 2 – planting, 3 – active maintenance, and 4 – monitoring and adaptive management. The vast majority of currently active restoration zones have yet to proceed to phase 4, and establishment of stable native communities often takes many years after the completion of active restoration. Analyses of restoration outcomes in this report should thus be viewed as baselines for future research and potential inputs for adaptive management or project planning, rather than settled

assessments of success in individual areas, as restoration has yet to be “completed” in most areas covered.

In order to assess the impact of GSP restoration activities on bird communities, we compared mean abundance (the average number of individuals per survey) and mean annual species diversity (the number of species reported annually per point) among survey points located within 50 meters of restoration zones in each phase, using a Tukey test to ask if there is a significant difference in either value across points in different restoration phases (Figure 7). This analysis found no significant difference in either diversity or abundance between points that had or had not undergone restoration, or among differing levels of restoration point class. Although the data present some suggestive trends of decreasing abundance and

FIGURE 7

Mean species diversity across stations located within 50m of areas at different stages of progress in habitat restoration. Error bars report one standard error.



diversity in phase 3 or 4 zones, this variation is within mean standard error of phase 1 and 2 zones and is not significant ($p > 0.05$) in Tukey tests. It should be noted that this test as currently employed has limited power because very few of the zones have proceeded beyond phase 3, and most are in phases 1 or 2. Apparent lower species diversity in phase 4 zones, for example, is largely driven by just two low-diversity points.

For all NBP survey points adjacent to GSP zones and with data series extending before the initiation of restoration activities, we also compared mean abundance and annual species diversity before and after the initiation of restoration work and used a paired t-test to assess the significance of any difference found. This procedure was repeated across species groups and focal species (Figure 8, Table 4).

We found that mean abundance declined for all species and groups assessed after the onset of restoration activities, though only total bird abundance, riparian bird abundance, and human-associated bird abundance showed significant differences in a t-test. The decline in human-associated birds explained roughly three-quarters of the decline in total bird abundance, which should be viewed as a cautious success for restoration, as these species are already abundant in surrounding urban habitats and often outcompete native species in heavily disturbed areas. The significant decline in riparian birds is a more worrying sign for the impact of restoration projects on bird communities, but this drop was nearly entirely explained by the significant decline in Song Sparrow populations – likely a reflection of the Song Sparrow’s success in living with Himalayan Blackberry, one of the most common invasive species removed during phase-1 restoration activities.

FIGURE 8

Mean species diversity across stations located within 50m of areas at different stages of progress in habitat restoration. Error bars report one standard error.





TABLE 4

Comparisons of measures of avian diversity and abundance before and after initiation of GSP habitat restoration work. Significantly different measures are in bold.

	PRE-RESTORATION MEAN	ONGOING RESTORATION MEAN	DIFFERENCE	<i>t</i>	<i>p</i>
Per-survey Species Diversity	5.126	5.307	0.181	1.883	0.063
Annual Species Diversity	6.998	7.966	0.968	1.757	0.082
Total Birds	17.123	14.898	-2.225	3.913	1.75E-4
Riparian Birds	2.175	2.018	-0.157	2.368	0.020
Invasive Species	9.499	7.524	-1.975	1.314	0.195
Warblers	2.000	1.908	-0.093	0.329	0.743
Woodpeckers	1.347	1.339	-0.008	0.164	0.870
Human-associated Birds	5.757	4.090	-1.667	4.340	3.65E-5
Wilson's Warbler	1.307	1.282	-0.025	0.230	0.819
Savannah Sparrow	2.114	1.934	-0.180	1.150	0.262
Anna's Hummingbird	1.278	1.244	-0.033	0.842	0.413
White-crowned Sparrow	2.066	1.732	-0.334	1.659	0.107
Orange-crowned Warbler	1.396	1.255	-0.141	1.273	0.211
Golden-crowned Kinglet	3.900	3.379	-0.521	1.723	0.089
Song Sparrow	2.093	1.923	-0.170	2.695	8.39E-3



© DOUG SCHURMAN

White-crowned Sparrow

Species diversity, measured both as mean annual diversity across survey points and as the average number of species reported per-survey per-point, increased on average by roughly one species per year after the initiation of restoration. Although this increase fell just short of statistical significance ($p=0.06$ per-survey, $p=0.08$ annual), the pattern is compelling and should be followed in future assessments of restoration impacts.

Overall, our assessment of the impact of GSP restoration activities on avian communities is cautiously positive. Observed declines in total bird counts (roughly 2 fewer birds per survey) are explained mostly by declines in counts of human-associated species, suggesting that restoration activities are, as intended, returning habitats to a more “natural” state less conducive to occupation by common urban birds. The consistent pattern of decline in abundance across species groups and focal species is somewhat worrying,

but likely reflects the ongoing disturbance caused by active work on a site as well as the time lag between establishment of native habitats in a restored area and establishment of bird populations using that habitat. Because most GSP restoration projects in the NBP study area were started in 2007 or later, very few zones have “completed” restoration. Thus our assessment provides a snapshot of the impact of active restoration on bird communities during a transitional phase in habitat quality. As more restoration zones are completed, we expect that patterns of increasing species diversity will continue, while patterns of declining abundance among non-human-associated species will level off; however, continued long-term monitoring will be necessary to assess these trends and make concrete recommendations for future restoration activity planning.

Left: Neighborhood Bird Project observations are recorded on data sheets, which are then entered into a database for later analysis. **Right:** A group of NBP volunteers identify birds using binoculars while surveying at Magnuson Park.



Magnuson Park Wetlands Restoration

From 2008 to 2011, Seattle City Parks undertook a large-scale habitat restoration project in Magnuson Park to remove invasive plants and hugely expand a complex of wetlands on the southern half of the park. This habitat restoration effort was much larger in scale than typical GSP sites assessed earlier in this report, encompassing an area over 14 acres and costing over \$3 million. Because NBP data collection in Magnuson Park began prior to the restoration and continued both during and post-construction, data from this site allows us to view how bird communities respond to restoration projects both during and after heavy construction. Because construction activities and changed topographies required some survey points to move, these data should be viewed as somewhat less conclusive than those from other NBP survey points, but the patterns observed are instructive and can help inform our view of how avian communities may respond to the end of work on the many smaller GSP restoration sites assessed here.

Counts of riparian birds were relatively constant from the beginning of data collection in 1998 through 2006, when

they experienced a slight decline. This decline persisted through the end of active construction in 2011, when riparian bird counts rebounded to roughly 50% above their pre-restoration baseline. Since 2011, frequency of riparian birds has been higher than in any year prior to the initiation of restoration. Meanwhile, abundance of human-associated species and invasive species has declined consistently since the beginning of data collection, reaching a minimum during the active construction phase and since maintaining relatively constant levels (though note that early results from 2014 surveys suggest a rebound in populations; Figure 9).

These patterns are encouraging early news for the success of this large project in increasing the abundance and diversity of native wildlife in Magnuson Park. The data also align well with standard expectations of progress in restoration projects, in which the highly visible short-term costs of large-scale construction to local wildlife are balanced by a long-term increase in abundance and diversity. Indeed, the speed with which riparian bird counts rebounded following construction – riparian bird frequency reached its maximum



A citizen science volunteer points out waterfowl which will be recorded at part of the Neighborhood Bird Project at Golden Gardens Park.



© DOUG SCHURMAN

Western Grebe


recorded level in the same year that heavy construction ended on the site – is surprising, and suggests that some of this increase in local abundance is the result of shifting populations from surrounding lower-quality habitats outside the park rather than an increase in the absolute number of riparian birds in the region, though an increase in available habitat should increase regional populations over the medium and long-term.

Viewing the observed trends in bird abundance and diversity around the smaller GSP restoration sites in the context of the Magnuson Park data, an optimistic interpretation would suggest that the observed declines in abundance across species near GSP sites are temporary and will be replaced by higher counts once restoration work is complete and sites are allowed time undisturbed for wildlife to discover the new habitats. The trends observed here also point to the critical role of long-term data collection in assessing the impact of restoration activities. In the case of Magnuson Park, this assessment of the response of avian communities was possible only because NBP’s volunteer surveys in the pre-restoration years had established a baseline level of bird abundance and diversity against which to compare the mid- and post-construction figures. NBP surveys to date have provided a similar baseline for many of the smaller GSP restoration sites and some early figures for in-progress sites are analyzed here, but assessment of overall restoration impacts will require continued data collection both through the active work phase (currently in progress in nearly all zones assessed) and after the completion of active work.


FIGURE 9

Species group abundance in Magnuson Park. Active constructions on the wetlands ran from 2008 to early 2011.

Mean Per-survey Abundance – **Magnuson Park**
INVASIVE SPECIES



Mean Per-survey Abundance – **Magnuson Park**
RIPARIAN SPECIES





© DOUG SCHURMAN

Song Sparrow



© DOUG SCHURMAN

Cedar Waxwing



© DOUG SCHURMAN

Black-capped Chickadee



© DOUG SCHURMAN

Red Crossbill



Bottom left: Many of the surveys for the Neighborhood Bird Project rely on “birding-by-ear” – identifying birds by their call only. A volunteer at Golden Gardens Park identifies birds by listening for calls.

Recommendations and Conclusions

As summarized in this report, the Neighborhood Bird Project has been successful in recording broad-scale trends in avian diversity and abundance in Seattle City Parks over its 17-year lifespan. Fueled by the efforts of over 330 dedicated volunteers, the program has provided an all-too-rare opportunity for community members to contribute meaningfully to science-based conservation and restoration projects in their own neighborhoods. With the growing interest in habitat restoration and its near-ubiquity across managed parklands in the Seattle area, long-term monitoring efforts like the NBP are also the most cost-effective way to gather the data necessary to make informed decisions about the management of some of our most heavily used public lands.

In order to maintain the integrity of the existing NBP dataset and to maximize its utility in future analyses, several modest improvements to survey methodology and design should be considered. First, although a partial distance-sampling protocol limiting observations to a 50m radius is included in the current NBP protocol, additional training or field protocols designed to ensure that surveys are limited to recording birds within 50m of an observation point should be implemented, to maintain the collection of high quality data over time by volunteers. The simplest measure available here would be to place flagging or otherwise visibly mark objects 50m from each observation point to give surveyors a frame of reference. Full distance sampling – the standard approach for professional avian point counts – involves recording the distance and direction to each individual bird recorded, but given the lack of this data for previous years and the difficulty of correctly locating and estimating distance to a bird detected only by sound without prior training, we do not recommend adopting this approach at this time.

Survey teams should also make every effort to avoid moving survey point locations, and should provide any new or moved point locations with new names rather than reusing the old ones. Although most survey points have remained in a single location throughout the period of data collection, construction, changes in topography, and changes in personnel over the years have all occasionally resulted in a point shifting location or being retired. Maintaining survey points in their historic locations for as long as possible will maximize the comparability of data across years, and keeping accurate records of locations is crucial to drawing any conclusions as to differences in bird community traits across landscapes.

Finally, the question of equality of effort between different surveyors is a constant worry both in volunteer and professionally conducted bird surveys. People with differing



Pileated Woodpecker

Bottom right: Citizen science volunteer using binoculars to identify and count birds for the Neighborhood Bird Project at Magnuson Park.

levels of familiarity with local birdsong, auditory capacity, visual acuity, and experience as a point counter can record very different numbers of birds in a given area, and over time these differences in surveyor ability can skew interpretation of the data. Although guaranteeing complete equality of ability between survey teams will never be possible for volunteer programs like the NBP, park leaders should do their best to ensure that all survey teams working on a given day are of roughly comparable ability. The most practical way to implement this recommendation is to ensure that each survey team has at least one member capable of birding by ear and identifying nearly all the birdsong heard during a point count on every survey.

Volunteers conduct surveys in all weather, here looking across the Puget Sound for waterfowl towards Bainbridge Island during a survey at Discovery Park.



© TIM BOYER

Turning to the trends in avian abundance and diversity documented in this report, we find grounds for cautious optimism as to the status of avian communities in Seattle City Parks. Both invasive species and human-associated species show long-term declines in abundance across all parks surveyed. Meanwhile, riparian birds, woodpeckers, and warblers - all groups that do well in native vegetation and are relatively scarce in the surrounding urban environments – are either increasing or holding steady in average abundance. Species-specific trends such as the marked decline in Savannah Sparrow abundance in Discovery Park point to the continued need for monitoring and adaptive management across the parks, and suggest that NBP data may provide useful information for land managers seeking to balance the needs of recreation and wildlife in the parks.

NBP data has also provided useful measures of the impacts of habitat restoration projects on avian communities. In Magnuson Park, a 14-acre wetland restoration project appears to have resulted in a marked increase in riparian bird abundance and coincides with the continued decline in abundance of invasive and human-associated species – both positive signs. Early observations from the many GSP restoration zones covering most of the parks included in the NBP dataset are more equivocal – diversity is slightly up, while abundance is down across the board. These declines in abundance may represent temporary impacts from active construction as were documented at Magnuson Park from 2008-2011, but long-term monitoring of GSP sites post-restoration will be necessary to draw better conclusions. The increases in species diversity, meanwhile, suggest that restoration has been modestly successful at introducing new habitat diversity to our parks. The trend should be watched carefully in the future.



Mew Gull




Belted Kingfisher



© TIM BOYER

APPENDIX

Golden Gardens NBP/GSP Zones




Carkeek Park NBP/GSP Zones



Discovery Park NBP/GSP Zones



Magnuson Park NBP/GSP Zones



Acknowledgements

This analysis was generously funded by the Sustainable Path Foundation who have a long history of supporting science based projects in the Puget Sound region.

The habitat data and work logs used throughout the analyses in this report were kindly provided by the Green Seattle Partnership and are based on thousands of hours of restoration work, often conducted by volunteers.

The Neighborhood Bird Project was conceived, developed and is managed by the Seattle Audubon Society. However, data collection relies exclusively on the hard work and expertise of more than 330 volunteer birdwatchers who have dedicated over 1300 hours to collecting data since the project's inception in 1994. To the right is a list of the (332 to date) volunteers who have taken part in the survey since 1994.



science for a healthy, sustainable future



Scientific Guidance

Eric Ward
Rob Faucett

Design

Lorie Ransom,
Tiny Whale
tinywhalecreative.com

Photography

Tim Boyer
timboyerphotography.com
Doug Schurman
flickr.com/photos/seattlebirdman
Tom Sanders

Abby Larson
Adam Sedgley
Al Ferkovich
Alan Grenon
Alice Arnold
Allisa Carlson
Alyssa Sampson
Amy Stevenson
Andres Ortiz
Andrew McCormick
Ann Hursey
Ann Zavitkovsky
Anna Kopitov
Anna Martin
Anne Bell
Anne Chafee
Anne Jacobs
Anne McCaffrey
Arn Slettebak
Art Eash
Barbara Blair
Barbara Broderick
Barbara Deihl
Barbara Retzlaff
Barbara Schnabel
Barbara Watts
Barbara Webster
Becky Benton
Becky Galloway
Berl Nussbaum
Bernard Martell
Bert Dudley
Beth Etscheid
Bianca Liebhaber
Bill Bennett
Bill Bogue
Bill Byers
Bill Shumway
Bo McFadden
Bob Benton
Bob Cleland
Bobby Pearson
Brandon Mathey

Brenda Senturia
Brett McCallum
Brian Etscheid
Brid Nowlan
Bryony Angell
Carl Haynie
Carleen Zimmerman
Carmine Pascucci
Carmine Vincenzo
Carole Wales
Carolee Colter
Caroline Cappello
Cathy Nolan
Charlene Ferkovich
Charles Crow
Charles Kahle
Charlie Cook
Charlotte Byers
Cheryl McArthur
Cheryl Petterson
Chris Karrenberg
Christi Norman
Christine Albright
Christine Gallegos
Colene McKee
Collin Vassallo
Curtis Pearson
Cynthia Wilson
Dan Clark
Dan Harville
Daniel Melamed
Darby Langdon
Dasha Gudalewicz
Darren Curtis
Darwin Lonso
Dasha Gudalewicz
David Hepp
David Saunders
David Whitten
Deborah Dowd
Diana Sorus
Diane Doles

Donald Fiene
Donna Luce
Donna Schaeffer
Eileen Bryant
Elaine Boehmer
Elaine Chuang
Elaine Magnusson
Elissa Ostergaard
Elizabeth Addis
Elizabeth Orians
Elizabeth Vincent
Emily Carter
Emily Fales
Emily Mandelbaum
Eric Carlson
Eric Wagner
Eric Ward
Erica Clark
Esther Neeser
Ethan Collins
Etta Cosey
Evan Hodder
Ezra Parker
Felice Levy
Fiona Cohen
Frank Montgomery
Fred Rowley
Fredianne Gray
Gary Kelsberg
Gene Hunn
George Ritchotte
Georgia Britt
Georgia Conti
Giles Pettifor
Gilia Angell
Noel Angell
Gina White
Gordon Orians
Gretchen Fetscher
Gwynneth Harris
Helen Crawford
Helen Gilbert
Helen Murphy



How you can help!

Please consider donating to Seattle Audubon to ensure the continuation of projects like these. More details at www.seattleaudubon.org.

If you're interested in joining one of the avian survey teams, contact the Science Manager at science@seattleaudubon.org and learn more about the project at www.seattleaudubon.org.

Helen Ross
 Helle Bielefeldt-Ohmann
 Henry Noble
 Herb Curl
 Jack Pauw
 Jackson Barnes
 James Nichols
 Jan Bragg
 Jan Harville
 Jan Lewinsohn
 Jane Hedberg
 Jane Johnson
 Janice Wagner
 Jay Donald Ostrow
 Jeanelle Richardson
 Jeannie Murphy-Ouellette
 Jeff Bryant
 Jeff Nystuen
 Jen Kunitsugu
 Jennifer Kauffman
 Jennifer McDonald
 Jennifer Riker
 Jenny Rapuzzi
 Jessica Kindred
 Jessica Piasecke
 Jessica Simmons
 Jim Thomas
 Joan Ostendorf
 Joe Anderton
 Joe Miles
 Joe Sweeney
 John Murtfeldt
 John Pauw
 John Tubbs
 Johnathan Britell
 JoLinda Finne
 Jon Woodard
 Joyce Moty
 Julia Allen
 Julia Bent
 Julie Monahan

Justine Busse
 Karen Adair
 Karen Stephens
 Kathryn Milburn
 Kathy Pstrak
 Kathy Slettebak
 Katie Barndt
 Ken Penner
 Kent Koprowicz
 Kent Slaven
 Kevin Castle
 Kimberly McDonald
 Kristin Johnson-Waggoner
 Kristin Marshall
 Laura Appell
 Laura Lipton
 Laurie Ann Dudley
 Lee Barnes
 Len Mandelbaum
 Libby Fiene
 Linda Carpender
 Linda Murtfeldt
 Linda Oswald
 Linda Phillips
 Linda Totle
 Lindsey Edwards
 Lisa Moore
 Liz Bell
 Lois Johnson
 Lorraine Hartmann
 Louise Martell
 Lynne Smith
 Maeve Lambert
 Mamie Bolender
 Marcia Kamin
 Marcia Stone
 Marianne Baker
 Marilyn Sandall
 Marilyn Sandall
 Marina Skumanich
 Mark Johnson
 Mark Salvadalena
 Mark Wolff

Martha Davis
 Martha Nester
 Martha Taylor
 Marti Louthier
 Marty Farrimond
 Marvin Breece
 Mary Ann Soltész
 Mary Anne Thorbeck
 Mary Roberts
 Mary Ullrich
 Mary Vincent
 Matt Bartels
 Matt Jensen
 Maureen McKelvey
 Maureen Traxler
 Maurie Kirschner
 Mayumi Tsuru
 Merrill Davison
 Mia Spangenberg
 Michael Carsiotis
 Michael Fleming
 Michael Witter
 Michele Herzberg
 Mickey Riley
 Mike Dermond
 Mike Freund
 Mira Lamb
 Miriam Gray
 Monya Noelke
 Nancy Edmondson
 Nancy Pearson
 Nate Starling
 Nathan Burkemoore
 Nathan Smith
 Neil Zimmerman
 Pam Cahn
 Pat Bredouw
 Pat Johnson
 Pat Little
 Patricia Hughes
 Patricia North
 Patricia Sanders
 Patti Brandt


Paul Meijer
 Paul Webster
 Paula Crockett
 Penny Bolton
 Penny Rose
 Peter Dunwiddie
 Peter Mann
 Philip Magasich
 Phyllis Moss
 Polly Radebaugh
 Rachel Lawson
 Raelene Gold
 Randolph Schnabel
 Randy Schnabel
 Rebecca Cahall
 Rebecca Evans
 Rene Koval-Huenuqueo
 Richard Youel
 Rita Condon
 Rob Faucett
 Robert Schmidt
 Roberta Roberts
 Roger Olstad
 Roland Kilcher
 Ron Leamon
 Ronald Paige
 Ruby Burkemoore
 Russ Kurtz
 Ruth Mynar
 Ruth Taylor
 Sage Gerow Miller
 Sam Woods
 Sarah Peden
 Sarah Safranek
 Scott Berglund
 Scott Hoskin
 Scott Ramos
 Shantel Gnewuch
 Sharon Aller
 Sharon Ellard
 Shiva Parameswaran
 Spencer Hildie
 Stephanie Burkemoore

Stephanie Durman
 Stephen Cass
 Steve Dang
 Steve Elston
 Steve Gerstle
 Sue Yates
 Susie Stillman
 Suzanne Hunter
 Tara Marino
 Ted Smith
 Temma Pistrang
 Teri Martine
 Terry Farris
 Terry Sisson
 Tiffany Linbo
 Tina Cohen
 Tom Weir
 Toni Potter
 Tor Linbo
 Tracee Geernaert
 Travis Keay
 Trisha Tubbs
 V. Linda Tofle
 Virginia Bound
 Wallis Bolz
 Wendy Croker
 Wendy Walsh
 William Denzel
 William Shumway
 Woody Wheeler
 Zoa Shumway

We attempted to capture everyone's name who has contributed in some way to the success of the Neighborhood Bird Project, if however, we've not listed your name here, we apologize and would love to hear from you.

NBP All Stars

Number of species observed by each volunteer surveyor from 1997-2014, limited to those who have observed 60+ species.



Recommended Citation

Batthey, C.J. and T. Ross. 2014. Impacts of Habitat Restoration and the Status of Avian Communities in Seattle City Parks. A Technical Report by the Seattle Audubon Society.

About the Authors

C.J. Batthey

Doctoral candidate at the Burke Museum of Natural History

CJ's main area of research involves systematics of new-world birds and the genetic control of migration in North American hummingbirds.

Toby Ross

Science Manager, Seattle Audubon

Toby manages and coordinates Seattle Audubon's three major Citizen Science projects, one of which is the Neighborhood Bird Project.

Front cover:

Golden-crowned Sparrow
©Doug Schurman



SEATTLE AUDUBON SOCIETY was founded in 1916 and is the oldest environmental organization in the state of Washington. Our mission is to cultivate and lead a community that values and protects birds and the natural environment. With approximately 5,000 members, we are one of the largest and most active Audubon chapters in the country. While legally separate organizations, we work closely with the 25 other local Audubon chapters in Washington state and with the state office of the National Audubon Society. Our program work seeks to connect people to birds and nature through environmental education, conservation advocacy, and citizen science projects. Volunteers are at the core of Seattle Audubon, with more than 600 dedicated, talented individuals contributing over 30,000 hours of service every year. A professional staff of 12 provides expertise and coordination for our program activities, working closely with our members and partner organizations.



BURKE MUSEUM OF NATURAL HISTORY AND CULTURE was founded in 1885. It is the oldest public museum in Washington and is a research- and collections-based museum with 16 million objects in its collections. The Burke Museum's Ornithology Collections exist primarily to inspire discovery. They generate new questions and provide data for testing ideas. The collections include: 41,000 study skins; 26,000 spread wings (the largest such collection in the world); 17,700 bird skeletons; 3,100 egg sets; and 26,000 avian tissues (the world's second largest collection). These bird specimens are used for teaching, research, and art.

